Brain glioma results by oncothermia

Prof. Dr. Andras Szasz^{1.2}

(1) Department of biotechnics, St. Istvan University, Budapest, Hungary (<u>Szasz, Andras@gek.szie.hu</u>) (2)
Oncotherm Group, Germany

Brain glioma results by oncothermia

Prof. Dr. Andras Szasz^{1,2}

(1) Department of Biotechnics, St. Istvan University, Budapest, Hungary (Szasz. Andras@gek. szie.hu) (2) Oncotherm Group, Germany

Objective: None of the established state-of-the-art treatments in malignant primary brain tumors, especially in glioblastoma multiform (GBM), could show effective or commonly accepted curative potential until today. The editorial question of JAMA [1] in 2005 is actual even now: "Where to go from here?" Our objective to show a feasible way to go, summarizing the results obtained till now made by modulated electro- hyperthermia (oncothermia) in various clinics in EU.

Method: Data are collected from GBM and anaplastic astrocytoma (AA) published observational studies. The method is transcranially applied modulated RF-current capacitive coupled at 13.56 MHz carrier frequency, (Oncotherm, EHY2000+) described in details elsewhere [2]. The applied protocol was unified step-up heating, 40-150 W RF-power with water-bolus cooling. Treatment is applied in combination with chemo- and/or radio-therapy or used as monotherapy if the conventional therapies fall.

Results: On brain-gliomas, the Groenemeyer Institute (Bochum Germany) was active [3], [4], [5], [6], [7], [8], [9], together with the BioMed Clinic (Bad Bergzabern, Germany), [10], [11], [12], [13], [14]; as well as the Empoli University had publication, [15]. Presently two German Universities (Regensburg [Prof. Bogdahn] and Heidelberg [Prof. Wick] are working on prospective clinical study on brain gliomas with hyperthermia. A dose escalating Phase I clinical study was made on Neurology Clinic, University of Regensburg, showing the safe process in the brain as well, [16]. The safe treatment could be shown by spectacularly documented near-eye cases, when the tumor disappeared by the oncothermia treatment, while the eye remained unhurt, intact from the treatment [17]. A well designed Phase I study is shown in other paper [18]. To see more evidences we show the retrospective data of independent clinics [19], having the same oncothermia protocol. The data are well correspond to each other and significantly higher than the data of the large international databases.

<u>Conclusion</u>: The results are strongly indicating the feasibility and the benefit of the oncothermia showing a valid treatment potential and safe application. Oncothermia is a potential way to escape from the present impasse situation and treat brain gliomas successfully. Performing prospective, randomized clinical trials in the future is mandatory.

References:

- $[1] \qquad \text{Fisher PG, Buffler PA: Malignant gliomas in 2005: where to GO from here? JAMA 293:615-617, 2005}$
- [2] Szasz A (2006) Physical background and technical realization of hyperthermia In: Baronzio GF, Hager ED (eds) Locoregional Radiofrequency- Perfusional- and Wholebody- Hyperthermia in Cancer Treatment: New clinical aspects, Ch 3, Springer Science Eurekah com, pp 27-59 http://www.springer.com/west/home/biomed/cancer/SGWID=4-125-22150459762-0&detailsPage=ppmmedia%7CotherBooks&seqNo=0&CIPageCounter=CI_MORE_BOOKS_BY_AUTHOR0
- [3] Sahinbas H, Baier Je, Groenemeyer DHW, Boecher E, Szasz A (2006) Retrospective clinical study for advanced brain-gliomas by adjuvant oncothermia (electro-hyperthermia) treatment www gimt-online de/uploads/media/Therapieergebnisse_Giloma_Studie_01 pdf
- [4] Sahinbas H, Groenemeyer DHW, Boecher E, Szasz A (2006) Retrospective clinical study of adjuvant electro-hyperthermia treatment for advanced brain-gliomas Deutche Zeitschrifts fuer Onkologie 39:154-160 http://www.thieme-connect.com/ejournals/abstract/dzo/doi/10 1055/s-2007-986020
- [5] Szasz A, Sahinbas H, Dani A (2004) Electro- hyperthermia for anaplastic astrocytoma and gliobastoma multiforme ICACT, Paris, 9-12 February 2004
- [6] Sahinbas H (2004) EHT bei Kindern mit Hirntumoren und nicht-invasive Messverfahren am beispiel von Hirntumoren Symposium Hyperthermie, Cologne, 15-16 October 2004
- [7] Sahinbas H, Grönemeyer D (2005) Oncothermia: An effective treatment for advanced gliomas Deutsche Kongress im Komplementare Medizine, Jun 2005, Schortens
- [8] Sahinbas H (2004) Deep RF hyperthermia treatment of advanced gliomas Oncology Conference, Basel
- [9] Sahinbas H, Grönemeyer D, Boecher E, Szasz A (2004) Hyperthermia treatment of advanced relapsed gliomas and astrocytoma The 9th International Congress on hyperthermic oncology, ICHO, St Louis, Missuri, 24-27 April 2004
- [10] Hager ED, Dziambor H, App EM, Popa C, Popa O, Hertlein M (2003) The treatment of patients with high-grade malignant gliomas with RF- hyperthermia Proc ASCO 22:118, #47;Proc Am Soc Clin Oncol 22: 2003 http://www.asco.org/portal/site/ASCO/menuitem_34d60f5624ba07fd506fe310ee37a01d/?vgnextoid=76f8201eb61a7010VgnVCM100000ed730ad1RCRD&vmview=abst_detail_view&confID=23&abstract D=102803
- [11] Hager ED, Sahinbas H, Groenemeyer DH, Migeod F (2008) Prospective phase II trial for recurrent high-grade malignant gliomas with capacitive coupled low radiofrequency (LRF) deep hyperthermia ASCO, Journal of Clinical Oncology, Annual Meeting Proceedings (Post-Meeting Edition) 26:2047 http://meeting ascopubs org/cgi/content/abstract/26/15_suppl/2047?maxtoshow=&HITS=10&hits=10&RESULTFORMAT=&fulltext=Hager&searchid=1&FIRS TINDEX=0&volume=26&issue=15_suppl&resourcetype=HWCIT
- [12] Hager ED (2004) Response and survival of patients with gliomas grade III/IV treated with RF capacitive-coupled hyperthermia ICHO Congress, St. Louis, USA
- [13] Hager ED (2004) Clinical Response and Overall Survival of Patients with Recurrent Gliomas Grade III/IV Treated with RF Deep Hyperthermia An Update ICHS Conference, Shenzhen, China
- [14] Fiorentini G, Giovanis P, Rossi S, Dentico P, Paola R, Turrisi G, Bernardeschi P (2006) A phase II clinical study on relapsed malignant gliomas treated with electro-hyperthermia, In Vivo 20:721-724 http://www.ncbi.nlm.nih.gov/sites/entrez?cmd=Retrieve&db=PubMed&list_uids=17203754&dopt=Citation
- [15] Fiorentini G, Giovanis P, Rossi S, Dentico P, Paola R, Turrisi G, Bernardeschi P: A phase II clinical study on relapsed malignant gliomas treated with electrohyperthermia, In Vivo 20:721-724, 2006
- [16] Wismeth C, Hirschmann B, Pascher C, Dudel C, Szasz A, Ogdahn U, Hau P (2009) Transcranial electro-hyperthermia combined with alkylating chemotherapy in patients with relapsing high-grade gliomas – Phase I clinical results Expanding the Frontiers of Thermal Biology, Medicine and Physics Annual Meeting of Society of Thermal Medicine, Tucson, USA, 3-7 April 2009
- [17] Cases from Prof Dr H Renner (Nuemberg, Germany) and Prof Dr A Herzog (Nidda, Germany) It could be sent by request for medical doctors
- [18] Wismeth C Dudel C Pascher C Ramm P Pietsch T Hirschmann B Reinert C Proescholdt M Rümmele P Schuierer G Bogdahn U HauP (2010) Transcranial electro-hyperthermia combined with alkylating chemotherapy in patients with relapsed high-grade gliomas Phase I clinical results, Journal of Neuro-Oncology 98:395-405 http://www.ncbi.nlm.nih.gov/sites/entrez
- [19] Szasz A (2009) Brain glioma results by oncothermia, a review Expanding the Frontiers of Thermal Biology, Medicine and Physics Annual Meeting of Society of Thermal Medicine, Tucson, USA, 3-7 April 2009